	T	
Farm Name		
Veterinary Practice Details		
•		
Type and number of Livestock covered	Cows	
by the plan	Heifers	
	Youngstock/ calves	
	Stock bulls	
	Store cattle	
	Finished cattle	
	(sheep)	
Names of persons who administer		
medicines		
		T
Plan completed by	Name:	Role on farm
	Vet Declaration: The information recorded within this	Signature:
	health plan, is, to the best of my	
	knowledge accurate and a true	
	reflection of practices on the farm.	Date:
	Review – Vet Declaration:	C:t
	The information recorded within this	Signature:
	health plan, is, to the best of my	
	knowledge accurate and a true	Date:
	reflection of practices on the farm.	Date.
	Review – Vet Declaration:	Signature:
	The information recorded within this	-
	health plan, is, to the best of my knowledge accurate and a true	
	reflection of practices on the farm.	Date:

Note: a separate sheep supplement is available. If the plan is used for Dairy, beef and sheep, the plans below must cover all cattle and sheep.

1. Biosecurity

Biosecurity measures must be farm-specific, relevant and proportionate to the health risks.

	Farm Specific Measures
Measures taken with incoming stock:	
Sourcing policy	
Ascertaining disease status	
Pre-movement/purchase information from seller	
Avoid / minimise mixing of stock (isolation period)	

Noi	ninated Isolation Facility/Area	
•	Location of nominated Isolation facility or area	
•	Actions required, if any, to make the facility available	
•	Means of cleaning and disinfecting after use	
Oth	er Measures:	
•	People	
•	Buildings	
•	Equipment	
•	Vehicles	
•	Farm pets (worming)	
*	Fencing	

2. Infectious Disease and Vaccinations

Disease	Status*	Test / Control Measure / Vaccine	Target Animals	Frequency/Timing
Johne's Disease				
(declaration to be				
made under NJMP)				
BVD				
Leptospirosis				
IBR				
ТВ				
Neospora				
Clostridial disease				

Use the blank rows to indicate other conditions that are relevant to the farm (eg calf pneumonia, Blue Tongue, salmonella, husk, ringworm)

3. Parasite Control

Parasite	Tests/ Control Measure / Product	Target Animals	Frequency/ Timing
Fluke			
Lungworm			
Intestinal Worms			
Lice			

^{*} Status: F - free, U - currently unknown, action required, V - vaccinated, T - being tested, M- being monitored, P - present, N - no action

Ticks		

Use the blank rows to include any other parasites that are relevant to the farm

4. Footcare and Lameness Management

Name of persons with responsibility for footcare and lameness management (mobility scoring and/or trimming)	Experience/ Qualifications

Mobility Scoring

Frequency	
Where recorded	
Action taken with	
Score 2 cows	
Score 3 cows	

Routine Measures taken for Prevention, Control and Treatment of Foot Problems (e.g. examination, trimming, footbathing)

Condition	Measure	When / How Often / Who	Products Used
Digital dermatitis	Control:		
	Treatment:		
Sole Ulcers	Control:		
	Treatment:		
Foul in the foot	Control:		
	Treatment:		
White line disease	Control:		
	Treatment:		
	Control:		
	Treatment:		
	Control:		
	Treatment:		

Use the blank rows to include other conditions that are relevant to the farm.

Describe method (s) of detection:

Routine preventative measures	Used?	Products used
Pre-dipping	Yes/ No	
Post-dipping	Yes/ No	
Cluster Spraying	Yes/ No	

Actions taken in treatment of clinical mastitis cases:

Condition	Measures Taken	Products Used	Route of administration	Withdrawal Period (milk and meat)
				Milk
				Meat
				Milk
				Meat
				Milk
				Meat
				Milk
				Meat

Please indicate details for any specific conditions being tackled.

6. Management of Drying Off

Procedures for drying off cows:

Group	Measures Taken	Products Used	Criteria for use/Thresholds	Withdrawal Period (milk and meat)
				Milk
				Meat
				Milk
				Meat
				Milk
				Meat
				Milk
				Meat

Please indicate if different actions are taken for different groups (e.g. high SCC cows)

7. Metabolic Disorders

Condition	Preventative Measures	Measure Taken If affected	Products Used
Hypomagnesaemia			
(Staggers)			
Milk fever			
Ketosis			
Displaced abomasum			

Use the blank rows to include other conditions that are relevant to the farm.

Jse of calving aid	Location	
	Rope cleaning (frequency, temp of water, cleaning agent)	
	Disinfectant used on aid and ropes	
	Calving aid used only when	
Average time calf left with row		
ohne's positive cow – action at calving		
Colostrum	Quantity	
	Quality	
	Timeframe	
	Method of feeding	
	Alternative source (frozen, powdered etc)	

9. Husbandry Procedures

Type and age of	Task	Person*	Method	Anaesthetic used
stock				
	Disbudding			
	Dehorning			
	Castration			

Training Records: I Identifying Treated Animals Describe the means used for identification (for all ages and species covered by the plan): Pain Relief – all pain relief must be recorded in the medicine records Procedure/injury/action Product used Rate Person responsible Disbudding/dehorning ameness Difficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service cessary, amend with any additional or farm-specific actions: I solate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	e the blank rows to include other routine operations supplied to the herd. Persons undertaking veterinary related tasks must be suitable, experienced and/or trained, details of which must be recorded Training Records. I dentifying Treated Animals Describe the means used for identification (for all ages and species covered by the plan): Pain Relief – all pain relief must be recorded in the medicine records Procedure/injury/action Product used Rate Person responsible Disbudding/dehorning ameness Difficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessory, amend with any additional or farm-specific actions: I isolate confirmed reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.		val of		
e the blank rows to include other routine operations supplied to the herd. Persons undertaking veterinary related tasks must be suitable, experienced and/or trained, details of which must be record Training Records. Identifying Treated Animals Describe the means used for identification (for all ages and species covered by the plan): Pain Relief – all pain relief must be recorded in the medicine records rocedure/injury/action Product used Rate Person responsible ilisbudding/dehorning ameness Difficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service sessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animals (e.g. with marker spray, tail tapes). Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	e the blank rows to include other routine operations supplied to the herd. Persons undertaking veterinary related tasks must be suitable, experienced and/or trained, details of which must be recorded Training Records. I dentifying Treated Animals Describe the means used for identification (for all ages and species covered by the plan): Pain Relief – all pain relief must be recorded in the medicine records rocedure/injury/action Product used Rate Person responsible disbudding/dehorning ameness Difficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) recessary, amend with any additional or farm-specific actions: I isolate confirmed reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.		•		
Persons undertaking veterinary related tasks must be suitable, experienced and/or trained, details of which must be record Training Records. I dentifying Treated Animals Describe the means used for identification (for all ages and species covered by the plan): Pain Relief – all pain relief must be recorded in the medicine records Procedure/Injury/action Product used Rate Person responsible Disbudding/dehorning ameness Difficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service cessary, amend with any additional or farm-specific actions: I Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	Persons undertaking veterinary related tasks must be suitable, experienced and/or trained, details of which must be recorded Training Records. I dentifying Treated Animals Describe the means used for identification (for all ages and species covered by the plan): Pain Relief – all pain relief must be recorded in the medicine records Procedure/injury/action Product used Rate Person responsible Person responsible Pisbudding/dehorning ameness Difficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessory, amend with any additional or farm-specific actions: I solate confirmed reactor animals (e.g. with marker spray, tail tapes). Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.	Freeze	branding		
Persons undertaking veterinary related tasks must be suitable, experienced and/or trained, details of which must be record Training Records. I dentifying Treated Animals Describe the means used for identification (for all ages and species covered by the plan): Pain Relief – all pain relief must be recorded in the medicine records Procedure/Injury/action Product used Rate Person responsible Disbudding/dehorning ameness Difficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service cessary, amend with any additional or farm-specific actions: I Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	Persons undertaking veterinary related tasks must be suitable, experienced and/or trained, details of which must be recorded Training Records. I dentifying Treated Animals Describe the means used for identification (for all ages and species covered by the plan): Pain Relief – all pain relief must be recorded in the medicine records Procedure/injury/action Product used Rate Person responsible Person responsible Pisbudding/dehorning ameness Difficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessory, amend with any additional or farm-specific actions: I solate confirmed reactor animals (e.g. with marker spray, tail tapes). Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.				
Persons undertaking veterinary related tasks must be suitable, experienced and/or trained, details of which must be record Training Records. I Identifying Treated Animals Describe the means used for identification (for all ages and species covered by the plan): Pain Relief – all pain relief must be recorded in the medicine records Procedure/injury/action Product used Rate Person responsible Disbudding/dehorning ameness Difficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service cessary, amend with any additional or farm-specific actions: I Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g., with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	Persons undertaking veterinary related tasks must be suitable, experienced and/or trained, details of which must be recorded Training Records. I dentifying Treated Animals Describe the means used for identification (for all ages and species covered by the plan): Pain Relief – all pain relief must be recorded in the medicine records Procedure/injury/action Product used Rate Person responsible Disbudding/dehorning ameness Difficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessory, amend with any additional or farm-specific actions: I solate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal for the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.				
Training Records. I. Identifying Treated Animals Describe the means used for identification (for all ages and species covered by the plan): Pain Relief – all pain relief must be recorded in the medicine records rocedure/injury/action Product used Rate Person responsible iisbudding/dehorning ameness iifficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd urggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service ressary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	Training Records. Identifying Treated Animals Describe the means used for identification (for all ages and species covered by the plan): Pain Relief – all pain relief must be recorded in the medicine records rocedure/injury/action Product used Rate Person responsible isbudding/dehorning ameness ifficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd uggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - ressorry, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal for the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.	e the blank rows to include o	ther routine operations supplied	I to the herd.	
Pain Relief – all pain relief must be recorded in the medicine records Procedure/injury/action Product used Rate Person responsible Disbudding/dehorning ameness Difficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	Describe the means used for identification (for all ages and species covered by the plan): Pain Relief – all pain relief must be recorded in the medicine records Procedure/injury/action Product used Rate Person responsible Disbudding/dehorning ameness Difficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animals(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.		ary related tasks must be suitable, o	experienced and/or trained, details of	of which must be recorded
Actions for Dealing with Confirmed TB Reactors in the Milking Herd Suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service accessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	Product used Rate Person responsible Disbudding/dehorning Ameness Difficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd Euggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) Coessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.			d species covered by the plan):	
Actions for Dealing with Confirmed TB Reactors in the Milking Herd Suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service accessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	Product used Rate Person responsible Disbudding/dehorning Ameness Difficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd Euggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) Coessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.				
Procedure/injury/action Product used Rate Person responsible Disbudding/dehorning Ameness Difficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	Procedure/injury/action Product used Rate Person responsible Disbudding/dehorning ameness Difficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.				
Actions for Dealing with Confirmed TB Reactors in the Milking Herd Suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service accessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	Actions for Dealing with Confirmed TB Reactors in the Milking Herd Suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.				
Procedure/injury/action Product used Rate Person responsible Disbudding/dehorning Ameness Difficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	Procedure/injury/action Product used Rate Person responsible Disbudding/dehorning ameness Difficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.				
Actions for Dealing with Confirmed TB Reactors in the Milking Herd Suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service accessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	Product used Rate Person responsible Disbudding/dehorning Ameness Difficult calving Actions for Dealing with Confirmed TB Reactors in the Milking Herd Euggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) Coessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.				
Actions for Dealing with Confirmed TB Reactors in the Milking Herd Suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	Actions for Dealing with Confirmed TB Reactors in the Milking Herd Suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.	. Pain Relief – all pain relief i	must be recorded in the medicin	ie records	
Actions for Dealing with Confirmed TB Reactors in the Milking Herd Suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	Actions for Dealing with Confirmed TB Reactors in the Milking Herd Suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.	Procedure/injury/action	Product used	Rate	Person
Actions for Dealing with Confirmed TB Reactors in the Milking Herd Suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	Actions for Dealing with Confirmed TB Reactors in the Milking Herd Suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.				responsible
Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.	isbudding/dehorning			
Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.	ameness			
Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Services cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	Actions for Dealing with Confirmed TB Reactors in the Milking Herd suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.				
 Suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. 	 Suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended. 	Difficult calving			
 Suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. 	 suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended. 				
 suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. 	 suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended. 				
 Suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. 	 Suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended. 				
 Suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. 	 suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended. 				
 Suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. 	 suggested policy is provided below (further detail can be found on the TBHub website and via the TB Advisory Service) - cessary, amend with any additional or farm-specific actions: Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended. 				
 Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. 	 Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended. 				
 Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. 	 Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended. 	Actions for Dealing with Co	nfirmed TB Reactors in the Milki	ing Herd	
 Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. 	 Isolate confirmed reactor animal(s) as soon as diagnosed by the veterinary surgeon pending their removal from the farm. Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended. 	_			
 Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. 	 Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended. 	suggested policy is provided l	below (further detail can be foun		the TB Advisory Service) -
 Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. 	 Clearly identify reactor animals (e.g. with marker spray, tail tapes). Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended. 	suggested policy is provided l	below (further detail can be foun		the TB Advisory Service) -
 Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. 	 Milk reactor animals last and thoroughly clean equipment after use. Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended. 	suggested policy is provided be cessary, amend with any add • Isolate confirmed rea	below (further detail can be foun litional or farm-specific actions:	d on the TBHub website and via t	
Discard milk from reactors completely - exclude from bulk tank and do not feed to calves.	 Discard milk from reactors completely - exclude from bulk tank and do not feed to calves. If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended. 	suggested policy is provided becessary, amend with any add Isolate confirmed rea	below (further detail can be foun ditional or farm-specific actions: actor animal(s) as soon as diagno	od on the TBHub website and via to	
	• If disposing of milk via direct land spreading, apply a grazing interval - minimum 6 months recommended.	 suggested policy is provided becessary, amend with any add Isolate confirmed reathe farm. Clearly identify reactor 	below (further detail can be foun litional or farm-specific actions: actor animal(s) as soon as diagno or animals (e.g. with marker spra	od on the TBHub website and via too seed by the veterinary surgeon posed by the veterinary surgeon posed.	
		 suggested policy is provided becessary, amend with any add Isolate confirmed reathe farm. Clearly identify reactor Milk reactor animals I 	below (further detail can be foun ditional or farm-specific actions: actor animal(s) as soon as diagnor animals (e.g. with marker spralast and thoroughly clean equipm	od on the TBHub website and via to esed by the veterinary surgeon p y, tail tapes). ment after use.	pending their removal fro
	ner actions taken:	 suggested policy is provided becessary, amend with any add Isolate confirmed reathe farm. Clearly identify reactor Milk reactor animals I Discard milk from rea 	below (further detail can be foun litional or farm-specific actions: actor animal(s) as soon as diagnor animals (e.g. with marker spraylast and thoroughly clean equipmentors completely - exclude from	osed by the veterinary surgeon posed by the veterinary surgeon posed by the veterinary surgeon posed to calve the contract of	ending their removal fro
HET ACTIONS LAKEN:		 Isolate confirmed reathe farm. Clearly identify reactor Milk reactor animals I Discard milk from rea If disposing of milk via 	below (further detail can be foun litional or farm-specific actions: actor animal(s) as soon as diagnor animals (e.g. with marker spraylast and thoroughly clean equipmentors completely - exclude from	osed by the veterinary surgeon posed by the veterinary surgeon posed by the veterinary surgeon posed to calve the contract of	ending their removal fro
		 suggested policy is provided becessary, amend with any add Isolate confirmed reather farm. Clearly identify reactor Milk reactor animals I Discard milk from rea 	below (further detail can be foun litional or farm-specific actions: actor animal(s) as soon as diagnor animals (e.g. with marker spraylast and thoroughly clean equipmentors completely - exclude from	osed by the veterinary surgeon posed by the veterinary surgeon posed by the veterinary surgeon posed to calve the contract of	ending their removal fro
		 Isolate confirmed reather farm. Clearly identify reactor Milk reactor animals I Discard milk from rea If disposing of milk via 	below (further detail can be foun litional or farm-specific actions: actor animal(s) as soon as diagnor animals (e.g. with marker spraylast and thoroughly clean equipmentors completely - exclude from	osed by the veterinary surgeon posed by the veterinary surgeon posed by the veterinary surgeon posed to calve the contract of	ending their removal fro

13.	Actions taken with Down When is vet notified? Euthanasia decision tree	ner Cows –	
14	Habbles/Charles valing		
14.	Hobbles/Shackles policy Where is the date/reason What frequency are chec When is vet notified Euthanasia decision tree		
	Euthanasia/ on-farm kill ent the methods used	ing d and names of who carries out euthanasia.	
Anima	l Туре	Method Used	Person who carries it out
Youngs	stock		
Adult s	stock		

Methods of Disposal:

16 Other Disease and Management Problems

Condition	Prevention/Treatment Measures (including products used and withdrawal periods)